ACCEPTED

VIA NOTIFICATION LABEL NOT REVIEWED

June 12, 2019

New York State Department of Environmental Conservation Division of Materials Management





Division of Materials Management Pesticide Product Registration R

KILLS: Ants, Bed Bugs, Cockroaches, Centipedes, Crickets, Flies, Fleas, Fruit Flies, Gnats, Mosquitoes, Moths, Scorpions, Silverfish, Spiders (Brown Recluse, Black Widow, others), Ticks, Wasps and Other Listed Insect Pests

Homes, Apartments, Hotels, Schools, Nursing Homes, Hospitals, Office Buildings, Warehouses, Commercial and Industrial Buildings and Other Listed Buildings and Structures

ACTIVE INGREDIENTS:

Etofenprox (CAS #80844-07-1)	16.2%
Piperonyl butoxide* (CAS #51-03-6)	
OTHER INGREDIENTS:	19.0%
TOTAL:	100.0%

* (Butylcarbityl)(6-propylpiperonyl) ether and related compounds
This formulation contains 1.4 lbs of etofenprox and 5.6 lbs of piperonyl butoxide per
gallon. End use dilution is 0.25% etofenprox.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panels for additional Precautionary Statements

EPA Reg. No. 2724-804 EPA Est. No. 2724-TX-1 300526593

NET CONTENTS: 1 PT (16 FL OZ) 473 ml

To the extent consistent with applicable, law, Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

www.zoecon.com

Actisol is a trademark of Environmental Delivery Systems, Inc. Micro-Injector is a registered trademark of Whitmire Micro-Ingen Research Laboratories, Inc.

Zoecon, Zoecon with design and Zenproxicare registered trademarks of Wellmark International. Central Garden & Pet and Central Garden & Pet with design are trademarks of Central Garden & Pet Company.

©2010-2016 Wellmark International.



Zenprox®



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS—CAUTION

Harmful if swallowed. Causes moderate eyel irritation. Avoid contact with eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Repeated exposure to etofenprox can cause skin irritation.

EPA Reg. No. 2724-804 EPA Est. No. 2724-TX-1

1 300520323



2-1/8"

PAGE 10 2-1/4"

FRONT COVER (Page 1)

FIRST AID

Call a poison control center or doctor immediately for treatment advice.

swallowed

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

General Directions: ZOËCON®! 'ZENPROX® EC (ZENPROX® EC) provides i effective kill of fleas, cockroaches and other listed pests and can be applied as an indoor broadcast, spot, void, or crack and crevice treatment using a low pressure sprayer, void injector or other equipment! (see Specific Directions) in homes, attics and crawlspaces, apartment buildings, hotels, schools, day cares, nursing! thomes, hospitals, office buildings, correctional facilities, commercial and industrial buildings, warehouses, nonrfood/non-feed handling establishments, modes of transportation, institutions. ZENPROX® EC provides up to 28 days! residual control of fleas and up to 14 days. residual on other listed pest species.

Apply ZENPROX® EC to areas where target pests hide, such as baseboards, corners, storage areas, closets, around water pipes, doors and windows, attics and eaves, behind and under refrigerators, cabinets, sinks, furnaces, and stoves, the

underside of shelves, drawers and similar areas. Pay particular attention to cracks and crevices.

Remove pets and birds and cover fish aquariums before spraying. Do not allow children or pets to contact treated surfaces until spray has dried. Do not apply to toys or stuffed animals or related articles. Do not apply this product in occupied patient rooms or in any rooms/areas, while occupied by patients, the elderly, or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, sports facilities, etc.) in the immediate area where occupants are present.

MIXING DIRECTIONS

ZENPROX® EC mixes easily with water (diluent) and may be applied with most application equipment such as Micro-Injector® or Actisol® applicators. For best results, partially fill sprayer with water (diluent), add the labeled amount of ZENPROX® EC and fill to the required amount of water (diluent). Diluted spray mixture may be stored overnight and used the following day, but mixture should be agitated well prior to use.

TZENPROX® EC in labeled dilutions is generally non-staining and should not stain surfaces that cannot be spotted or damaged by water alone, however in questionable or sensitive surfaces, apply a small amount of dilution in an inconspicuous location to evaluate. Do not apply where electrical short circuits might result. Do not enter or allow others to enter until spray has dried. Do not apply product in a way that will contact workers, other persons, or animals, either directly or through drift.

TANK-MIXING

ZENPROX® EC can be mixed with EPA-laregistered insect growth regulators in areas where these products are registered to be applied. Please check product-laspecific label directions before using and the more stringent label directions will apply.

STORAGE and **DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry, and ventilated area away from humans, pets, foodstuffs, and feed. Do not store near heat or open flame. Keep from freezing.

Pesticide Disposal: To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticided disposal program (often such programs are run by state or local governments or by industry).

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple, rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use on disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or incinerate, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

SPECIFIC INSTRUCTIONS Indoor Pests: Use ZENPRCX® EC as a proadcast (general surface), spot, void, or crack and crevice treatment to control pests inside buildings, structures, and modes of transport. This includes homes, attics and crawlspaces, apartment buildings, hotels, schools, day cares, nursing homes, hospitals, office buildings, in correctional facilities, commercial and industrial buildings, warehouses, non-food/non-feed handling establishments, rail cars, marine vessels, and institutions. ZENPROX® EC mixes easily with water (diluents) and may be applied with a low pressure sprayer for broadcast and spot treatments. A void injector or other application equipment that delivers low volume treatments, such as Micro-Injector® or Actisol™ applicators, may be used for void and crack and crevice treatments. Pests **Use Rate Specific Directions** Treatment 0.25% Etofenprox Apply the dilution uniformly over surfaces to be Broadcast 2 fl oz/gallon treated. Apply to carpets, floors, cabinets, cove base, Ants, Bed Hugs, Carpet beetles, Cockroaches, Confused flour (General Surface) of water and other areas where pests are located. beetles, Cehtipedes, Crickets, Darkling beetles, Dermestids, (diluent)/1,000 ft2 Earwigs, Firebrats, Flies, Fleas, Fruit flies, Gnats, Indian Treat entry points such as around water pipes, meal moths, Lesser grain borers, Millipedes, Mosquitoes, doors, windows, and eaves, Treat areas where pests 0.25% Etofenprox Moths, Pillbugs, Red flour beetles, Saw-toothed grain beetles, normally feed or hide such as haseboards, corners, 2 fl oz/gallon Scorpions, Silverfish, Sowbugs, Spiders (Brown recluse, Black Spot, Void or around water pipes, cracks and crevices. of water widow, others. Stink bugs, Ticks, Wasps Crack and Crevice (diluent)/1,000 ft2 Treat void areas with sufficient material to cover the void surfaces. Spot treat floorirugs, beneath furniture, in closets, and storage areas. Disassemble the bed frame and remove bedding. Do not apply product to bedding. Wash bedding and dry using medium to high heat to kill bed bugs. Apply to the bed frame, mattress and box spring. Spot, Void or Crack 0.25% Etofenprox Application to mattresses and boxisprings is intended and Crevice for 2 fl oz/gallon for treatment of seams, edges, and tufts only. Do not Bed Bugs Infesting Bed bugs of water saturate the mattress and box spring. Do not treat the Beds, Mattresses, (diluent)/1,000 ft² entire surface of the mattress dr box spring. Allow Bedrooms and spray to dry before using the mattress and/or box Sleeping Areas spring. Spot treat baseboards, behind pictures/wall hangings, rugs, and other articles in the bedroom or sleeping area. Repeat treatments to infested areas as necessary, but not less than 14 days after the initial treatment.

Zenprox® Ec

'ACTIVE INGREDIENTS:

Etofenprox (CAS #80	844-07-1)	16.2%
Piperonyl butoxide*	(CAS #51-03-6)	64.8%

OTHER INGREDIENTS: 19.0%

* (Butylcarbityl)(6-propylpiperonyl) ether and related compounds

This formulation contains 1.4 lbs of etofenprox and 5.6 lbs of piperonyl butoxide per gallon. End use dilution is 10.25% etofenprox.

KEEP OUT OF REACH OF CHILDREN CAUTION

Refer to attached leaflet for complete Directions for Use and Precautionary Statements

EPA Reg. No. 2724-804

IEPA Est. No. 2724-TX-1

. |Wellmark International

1501 East Woodfield Rd., 200W Schaumburg, Illinois 60173

300520323

NET CONTENTS: 1 PINT (16 FL 0Z) 473 ml